

CLINICAL PRACTICE

Long-QT Syndrome

Dan M. Roden, M.D.

This Journal feature begins with a case vignette highlighting a common clinical problem. Evidence supporting various strategies is then presented, followed by a review of formal guidelines, when they exist. The article ends with the author's clinical recommendations.

After the sudden death of a 13-year-old girl while she was playing basketball, her family comes to the clinic for medical evaluation (Fig. 1). Her parents' resting electrocardiograms (ECGs) are normal, but her 9-year-old sister's ECG shows an abnormally long QT interval. There is a history of recurrent syncope in female relatives of the maternal grandmother, but there is no family history of other sudden deaths, the sudden infant death syndrome, drowning, or death from a motor vehicle accident. How should these family members be further evaluated and treated?

THE CLINICAL PROBLEM

Sudden death in an otherwise healthy young person is an emotionally devastating event and unfortunately an all-too-common presentation of genetic arrhythmia syndromes. The most common of these syndromes is the long-QT syndrome, which is characterized by abnormal QT-interval prolongation on the surface ECG and an increased risk of sudden death, usually due to ventricular fibrillation. Physical stress and emotional stress are common triggers of syncope or sudden death in the long-QT syndrome; occasionally these events are triggered by loud noises or occur while the person is at rest.¹⁻³

Hundreds of mutations in 10 genes linked to the long-QT syndrome have been identified (Table 1; and Table A in the Supplementary Appendix, available with the full text of this article at www.nejm.org).^{9,10} Mutations in three genes, each encoding a cardiac ion channel that is important for ventricular repolarization, account for the vast majority of cases; the resulting genetic subtypes are called LQT1, LQT2, and LQT3 (Table 1). Most reported mutations are in coding regions, although non-coding mutations (resulting in the loss of allele expression) have also been described.¹¹⁻¹³ Most families with the condition have their own mutations, which are often termed "private" mutations; an emerging body of data suggests that the location of the altered amino acid (or acids) within the ion-channel proteins may affect the prognosis.^{14,15} Some clinical features such as QT morphologic characteristics, the response of the QT interval to exercise, triggers of arrhythmia, and the response to therapies vary according to the disease-associated gene. Disease-associated genes (described in Table A in the Supplementary Appendix) are rare, and generalizations regarding gene-specific presentations or responses to therapy are therefore less reliable than those in common subtypes.

Syncopal patients with the long-QT syndrome is generally attributed to the form of polymorphic ventricular tachycardia called torsades de pointes. The LQT3 form of the syndrome can also be associated with bradycardia, and slow heart rates may cause syncope in some patients. Death is usually due to ventricular fibrillation. Most cases are associated with the autosomal dominant form of the syndrome (i.e., the Romano-Ward syndrome), with striking variability in clinical phenotypes

From the Departments of Medicine and Pharmacology, Vanderbilt University School of Medicine, Nashville. Address reprint requests to Dr. Roden at the Departments of Medicine and Pharmacology, Oates Institute for Experimental Therapeutics, Vanderbilt University School of Medicine, 1285 Medical Research Building IV, Nashville, TN 37232-0575, or at dan.rodan@vanderbilt.edu.

N Engl J Med 2008;358:169-76.

Copyright © 2008 Massachusetts Medical Society.

Table 1. Common Forms of the Long-QT Syndrome.*

Variable	Genetic Subtype		
	LQT1	LQT2	LQT3
Disease-associated gene	<i>KCNQ1</i>	<i>KCNH2</i>	<i>SCN5A</i>
In vitro effect	Decreased I_{Ks}	Decreased I_{Kr}	Increased plateau I_{Na}
Setting of arrhythmia†	Emotional or physical stress, swimming, diving	Emotional or physical stress, sudden loud noise	Rest, sleep
Typical resting ECG‡	Broad T wave	Low-amplitude T wave with notching	Long isoelectric ST segment
ECG at onset of arrhythmia§	No pause	Pause	Not established
QT change with exercise	Failure to shorten	Normal	Supranormal
QT shortening with mexiletine¶	No	No	Yes
Clinical response to beta-blockers	Yes	Less than LQT1 response	Uncertain

* ECG denotes electrocardiogram, I_{Kr} the rapid component of the delayed rectifier current, I_{Ks} the slow component of the cardiac delayed rectifier current, and I_{Na} the cardiac sodium current.

† Data are from Schwartz et al.⁴

‡ These are typical patterns, but exceptions and variants are well recognized. Data are from Moss et al.⁵

§ Data are from Tan et al.⁶

¶ Data are from Schwartz et al.⁷

|| Data are from Priori et al.⁸

How should mutation carriers be counseled and treated?

STRATEGIES AND EVIDENCE

DIAGNOSIS

Common presentations of the long-QT syndrome are palpitations, presyncope, syncope, and cardiac arrest. In addition, asymptomatic persons may be evaluated because the diagnosis is established or suspected in a family member. The differential diagnosis includes common causes of syncope in the young; these causes range from benign conditions such as vasovagal syncope to serious genetic conditions such as hypertrophic cardiomyopathy and catecholaminergic polymorphic ventricular tachycardia.²⁶⁻²⁸ The history can help with the differential diagnosis and point toward specific subtypes of the long-QT syndrome. The physical examination and echocardiography (or magnetic resonance imaging, if performed) show no abnormalities in patients with this syndrome; thus, they are helpful only in ruling out other diagnoses.

An abnormal ECG obtained while the patient is at rest is the key to diagnosis. The QT interval, the surface ECG representation of ventricular repolarization, is affected by the patient's heart rate and sex; the upper limits of the QT interval corrected for the heart rate (the QTc) are below 460 msec for women and below 440 msec for

men. Yet establishing a diagnosis of the long-QT syndrome may not be straightforward. Physicians may misread the QT interval²⁹ or misdiagnose vasovagal syncope as the long-QT syndrome.²⁶ In addition, mutation carriers may have normal ECGs, although this is unusual in patients with symptoms. The Schwartz scoring system, which includes ECG features and personal and family history, has been widely used,³⁰ but it does not take into account genetic information.³¹ All persons with QT-interval prolongation should be screened for acquired causes such as hypocalcemia, hypothyroidism, and the use of drugs that can prolong the QT interval; these drugs include antiarrhythmic agents such as sotalol and dofetilide and noncardiovascular drugs such as haloperidol, methadone, and pentamidine. Most drugs that cause torsades de pointes block the rapid component of the delayed rectifier current (I_{Kr}), and previously unrecognized long-QT syndrome, of any subtype, can be identified in 5 to 20% of patients with drug-induced torsades de pointes.³²⁻³⁴

A detailed family history is essential. This evaluation should assess not only a history of sudden death, but also other deaths that may have been a manifestation of the long-QT syndrome, such as drowning, the sudden infant death syndrome,³⁵ and the death of a family member while he or she was driving (a potential manifestation of syncope).

Exercise testing can be useful to assess the

response of the QT interval (Fig. 1B and Table 1), but arrhythmias are very rare. The epinephrine challenge has been used to identify mutation carriers in families with the LQT1 mutation.^{22,36} Holter monitoring and electrophysiological testing are generally not useful.

GENETIC TESTING

The long-QT syndrome is a clinical diagnosis, but genetic testing may provide additional information. Clinical features such as triggers of syncope and specific QT morphologic attributes^{5,37} (Fig. 2) in patients in whom the clinical diagnosis has been made can suggest the affected gene in 70 to 90% of patients.^{38,39} Genetic testing for the common subtypes of the long-QT syndrome is now available commercially, and it can identify a mutation in 50 to 75% of probands in whom the di-

agnosis appears to be certain on clinical grounds. The lack of detection in the remainder of the probands is probably due to technical difficulties with genotyping,⁴⁰ noncoding variants,¹¹⁻¹³ or as yet unidentified disease-associated genes. Thus, a negative genetic test does not rule out the diagnosis. There is also the potential for false positive results, since detection of a previously undescribed mutation does not establish the diagnosis. Rather, further analysis (e.g., linkage within a family or in vitro studies) may be required to establish the functional significance of any given variant; rare DNA variants of little functional consequence are well recognized.

Genetic testing for the long-QT syndrome is most useful in two settings. First, when a clinical diagnosis is relatively certain, knowing the specific gene affected (or the site of the mutation within the gene)^{14,15} may clarify the prognosis and guide therapeutic choices. Second, in a family with an affected proband and a known genetic defect, the genotyping of family members can help rule out the diagnosis in some persons. However, a positive test identifies a family member as being a mutation carrier, even if he or she is asymptomatic, has a normal QT interval, and is unlikely ever to have an event; this identification may have attendant consequences for long-term therapies and screening in his or her own family. A positive genetic test also could be used in a discriminatory fashion (e.g., to withdraw health insurance), although the pending Genetic Information Nondiscrimination Act is meant to remedy this situation. Detailed genetic counseling is warranted before proceeding to this testing, particularly for asymptomatic persons for whom the option of not testing must also be recognized.

Genetic testing has not been evaluated in patients who present with a borderline QT interval, suspicious symptoms (e.g., syncope), and no relevant family history. In these patients, the incidence of false positive and false negative results and their implications for management remain unknown.

RISK STRATIFICATION

The most powerful predictor of risk is the QTc duration.^{20,41-43} In an analysis of 647 LQT1, LQT2, and LQT3 mutation carriers, the incidence of syncope or sudden death by 40 years of age in those with a QTc interval in the lowest quartile (<446 msec) was less than 20%, whereas it was




Figure 2. Electrocardiographic Patterns in the Three Common Forms of the Long-QT Syndrome.

The LQT1 form of the long-QT syndrome is associated with a broad T wave without a shortening of the QT interval due to exercise (as shown in Fig. 1C). LQT2 is associated with low-amplitude, often bifid, T waves. LQT3 is associated with a long isoelectric segment and a narrow-based, tall T wave. Although patterns may suggest a specific genotype of the long-QT syndrome,⁵ many variants have been described.³⁷

more than 70% among those in the highest quartile (>498 msec).²⁰ In the same study, features identifying patients at particularly high risk (>50% risk of an event before 40 years of age) included a QTc interval of more than 500 msec in carriers with LQT1 and LQT2 and male sex in carriers with LQT3; predictors of lower risk (<30% risk of an event before 40 years of age) included a QTc interval of less than 500 msec in carriers with LQT1 and male sex in carriers with LQT2. Although some of the subgroups were small and the effects of therapy were not controlled, these findings underscore the concept that risk is a continuum in this disease. Mutations that appear to be especially severe have been reported,^{14,44} although the numbers of affected patients and families are small.

THErapy

The lack of randomized trials of therapy in the long-QT syndrome reflects both the relative rarity of the disease and the heterogeneity in the type and severity of its clinical presentations. Data to guide management are generally from large registries and referral centers and thus are biased toward patients with severe disease.

Persons with a very low risk of sudden death (e.g., elderly mutation carriers with normal QT intervals) need not be treated, although even in these persons, it is prudent to avoid drugs known to prolong the QT interval. The major therapeutic options for the long-QT syndrome are beta-blockers and implantable cardioverter-defibrillators (ICDs).

The mainstay of therapy for the long-QT syndrome has been beta-blockade. Long-acting preparations such as nadolol and atenolol are usually used, and the efficacy of beta-blockade is assessed by blunting of the exercise heart rate (e.g., by >20%); beta-blockers do not substantially shorten the QT interval. Extensive observational data before¹⁻³ and after⁴² the identification of disease-associated genes have shown superior survival among symptomatic patients who received beta-blockers (or occasionally among those who underwent left stellate ganglionectomy)⁴⁵ as compared with those who did not. Observational data have suggested that syncope or sudden death is less likely during beta-blocker therapy among patients with the LQT1 subtype (reported event rate among treated patients, 10% by 40 years of age) than among those with LQT2 or LQT3⁸; these

findings are consistent with the adrenergic dependence of LQT1.

Sodium-channel blockers such as mexiletine and flecainide may normalize the QTc interval in patients with the LQT3 subtype,^{46,47} but they may also increase the risk of sudden death in patients with overlapping Brugada syndrome^{48,49}; their role as primary therapy in LQT3 thus remains uncertain. Identification of patients with both the LQT3 subtype and the Brugada syndrome may require further study (e.g., with the use of ECGs after monitored sodium-channel blocker challenge⁴⁸ or in vitro studies⁴⁹).

The use of ICDs is widely considered in patients at high risk for sudden death, including those with symptoms before puberty, those with very long QTc intervals (e.g., >500 msec), and those with recurrent syncope thought to be due to arrhythmias, despite adherence to an adequate beta-blocking regimen. Some data also suggest that patients with cardiac arrest rather than syncope as the presenting symptom are at higher risk for sudden death.⁵⁰ Because of the adrenergic component triggering arrhythmias in the long-QT syndrome, the use of an ICD may be complicated by multiple shocks.⁵¹

AREAS OF UNCERTAINTY

The mechanisms underlying the variable penetrance of long-QT syndrome mutations are unknown. The elucidation of these mechanisms might facilitate risk stratification.

Because syncope or death in the long-QT syndrome is often adrenergically mediated, restriction of patients' participation in athletic activities is generally recommended. It is not known whether this restriction should extend to patients with forms of the disease in which adrenergic stressors are not prominent.

The appropriate role of ICDs remains controversial. Available data indicate that the severity of symptoms in the proband does not predict severity in affected family members.⁴¹ A conservative approach is to reserve the use of ICDs for patients with symptoms that occur despite the use of beta-blocker therapy. A more aggressive approach is to implant an ICD in any affected person, including persons in whom the long-QT syndrome has been detected by means of family screening and in whom there is evidence of an increased risk of sudden death. An argument for

the liberal use of ICDs is their lifesaving potential in young, otherwise healthy patients. However, the associated costs and the observation that most patients who have mutations will never have cardiovascular events argue against this course.

Long-term follow-up studies involving patients receiving ICDs for primary prevention in this setting are needed to inform decision making. Further study may also identify genetic or sophisticated ECG markers that predict either increased or blunted risk in a mutation carrier, but it seems unlikely that any test will show that a carrier has no risk.

Finally, there is inevitably an intense wish on the part of families and physicians to avoid any risk after the death of a young mutation carrier. Thus, ICDs are likely to be used more as family screening becomes more common. ICDs are generally implanted in young patients who will have the device for decades. Advances that minimize even small risks associated with ICDs, such as component failure or infection, would be especially beneficial in younger patients.

GUIDELINES

The major cardiology and electrophysiology societies in the United States and Europe have jointly issued guidelines for the care of patients who are at risk for sudden death from cardiac causes.

These patients include those with the long-QT syndrome (Table 2).⁵²

CONCLUSIONS AND RECOMMENDATIONS

The evaluation of family members after the sudden death of a young person, as described in the vignette, should start with a detailed family history to elicit information about any other cases of sudden death, as well as deaths due to the sudden infant death syndrome, drowning, and motor vehicle accidents. Autopsy findings are usually normal in patients with fatal long-QT syndrome. If the proband has survived a cardiac arrest, imaging to rule out structural diseases is also normal. The key to diagnosis is the resting ECG; although a long QT interval suggests the syndrome, other causes of QT prolongation (e.g., hypocalcemia or hypothyroidism) should be ruled out. Occasionally, the diagnosis will become apparent only with provocation such as treadmill exercise or an epinephrine challenge. Families often come to clinical attention when a healthy young person has died, and grief counseling is integral to patient care in this setting.

Once the diagnosis of the long-QT syndrome has been established, further history, especially the circumstances surrounding the syncope or sudden death, and occasionally features of the ECG and genetic testing may assist in identifying

Table 2. Guidelines for Management of the Long-QT Syndrome.*

Recommendation	Level of Evidence†	Comment
No participation in competitive sports	I	Includes patients with the diagnosis established by means of genetic testing only
Beta-blockers	I	For patients who have QTc-interval prolongation (>460 msec in women and >440 msec in men)
Implantable cardioverter–defibrillator	IIa	For patients with a normal QTc interval
	I	For survivors of cardiac arrest
	IIa	For patients with syncope while receiving beta-blockers
	IIb	For primary prevention in patients with characteristics that suggest high risk; these include LQT2, LQT3, and QTc interval >500 msec‡

* Data are from the American College of Cardiology, the American Heart Association, and the European Society of Cardiology, in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. Guidelines are adapted from Zipes et al.⁵²

† Levels of evidence are as follows: I, conditions for which there is evidence or general agreement, or both, that a given procedure or treatment is beneficial, useful, and effective; II, conditions for which there is conflicting evidence or divergence of opinion, or both, about the usefulness and efficacy of a procedure or treatment; IIa, conditions for which the weight of evidence or opinion is in favor of usefulness and efficacy; and IIb, conditions for which the usefulness and efficacy are less well established by evidence or opinion.

‡ Other indicators of risk may include the specific site of mutation¹⁴ and the postpartum period.¹⁸

a subtype. Genetic testing is also useful to establish or rule out the diagnosis in a family member of a patient with a known genetic mutation.

Beta-blockade with a long-acting agent is the mainstay of therapy. The use of an ICD should be considered, particularly when features suggesting an unusually high risk of sudden death are present; these features include especially long QT intervals, the onset of syncope with sudden noise or at rest, and certain ECG patterns (Fig. 2). In the family described in the vignette, beta-blockade is indicated in the 9-year-old sister and the mother (who is 39 years of age and carries the LQT1 mutation), but it would not be recommend-

ed for the maternal grandfather. The use of an ICD may be considered in the 9-year-old girl because she has one high-risk feature (QTc interval >500 msec), but this approach remains controversial, reflecting the challenges in managing this condition.

Supported by grants from the U.S. Public Health Service (HL65962 and HL49989) and from the Fondation Leducq, Paris (05-CVD-01).

Dr. Roden reports receiving consulting fees from Sapphire Therapeutics, Atlas Venture Advisors, Pfizer, Avanir, Baker Brothers Advisors, CardioKine, and Eli Lilly and holding a patent on D85N as a predictive single-nucleotide polymorphism for drug-induced long-QT syndrome. No other potential conflict of interest relevant to this article was reported.

REFERENCES

- Moss AJ, McDonald J. Unilateral cervicothoracic sympathectomy for the treatment of long QT interval syndrome. *N Engl J Med* 1971;285:903-4.
- Schwartz PJ, Periti M, Malliani A. The long QT syndrome. *Am Heart J* 1975;89:378-90.
- Moss AJ, Schwartz PJ, Crampton RS, Locati E, Carleen E. The long QT syndrome: a prospective international study. *Circulation* 1985;71:17-21.
- Schwartz PJ, Priori SG, Spazzolini C, et al. Genotype-phenotype correlation in the long-QT syndrome: gene-specific triggers for life-threatening arrhythmias. *Circulation* 2001;103:89-95.
- Moss AJ, Zareba W, Benhorin J, et al. ECG T-wave patterns in genetically distinct forms of the hereditary long QT syndrome. *Circulation* 1995;92:2929-34.
- Tan HL, Bardai A, Shimizu W, et al. Genotype-specific onset of arrhythmias in congenital long-QT syndrome: possible therapy implications. *Circulation* 2006;114:2096-103.
- Schwartz PJ, Priori SG, Locati EH, et al. Long QT syndrome patients with mutations of the *SCN5A* and *HERG* genes have differential responses to Na⁺ channel blockade and to increases in heart rate: implications for gene-specific therapy. *Circulation* 1995;92:3381-6.
- Priori SG, Napolitano C, Schwartz PJ, et al. Association of long QT syndrome loci and cardiac events among patients treated with beta-blockers. *JAMA* 2004;292:1341-4.
- Moss AJ, Kass RS. Long QT syndrome: from channels to cardiac arrhythmias. *J Clin Invest* 2005;115:2018-24.
- Keating MT, Sanguinetti MC. Molecular and cellular mechanisms of cardiac arrhythmias. *Cell* 2001;104:569-80.
- Li H, Chen Q, Moss AJ, et al. New mutations in the *KVLQT1* potassium channel that cause long-QT syndrome. *Circulation* 1998;97:1264-9.
- Zhang L, Vincent GM, Baralle M, et al. An intronic mutation causes long QT syndrome. *J Am Coll Cardiol* 2004;44:1283-91.
- Koopmann TT, Alders M, Jongbloed RJ, et al. Long QT syndrome caused by a large duplication in the *KCNH2* (*HERG*) gene undetectable by current polymerase chain reaction-based exon-scanning methodologies. *Heart Rhythm* 2006;3:52-5.
- Moss AJ, Shimizu W, Wilde AAM, et al. Clinical aspects of type-1 long-QT syndrome by location, coding type, and biophysical function of mutations involving the *KCNQ1* gene. *Circulation* 2007;115:2481-9.
- Moss AJ, Zareba W, Kaufman ES, et al. Increased risk of arrhythmic events in long-QT syndrome with mutations in the pore region of the human ether-a-go-go-related gene potassium channel. *Circulation* 2002;105:794-9.
- Priori SG, Napolitano C, Schwartz PJ. Low penetrance in the long-QT syndrome: clinical impact. *Circulation* 1999;99:529-33.
- Imboden M, Swan H, Denjoy I, et al. Female predominance and transmission distortion in the long-QT syndrome. *N Engl J Med* 2006;355:2744-51.
- Seth R, Moss AJ, McNitt S, et al. Long QT syndrome and pregnancy. *J Am Coll Cardiol* 2007;49:1092-8.
- Zareba W, Moss AJ, Schwartz PJ, et al. Influence of the genotype on the clinical course of the long-QT syndrome. *N Engl J Med* 1998;339:960-5.
- Priori SG, Schwartz PJ, Napolitano C, et al. Risk stratification in the long-QT syndrome. *N Engl J Med* 2003;348:1866-74.
- Wang Q, Curran ME, Splawski I, et al. Positional cloning of a novel potassium channel gene: *KVLQT1* mutations cause cardiac arrhythmias. *Nat Genet* 1996;12:17-23.
- Vyas H, Hejlik J, Ackerman MJ. Epinephrine QT stress testing in the evaluation of congenital long-QT syndrome: diagnostic accuracy of the paradoxical QT response. *Circulation* 2006;113:1385-92.
- Wilde AA, Jongbloed RJ, Doevendans PA, et al. Auditory stimuli as a trigger for arrhythmic events differentiate *HERG*-related (*LQTS2*) patients from *KVLQT1*-related patients (*LQTS1*). *J Am Coll Cardiol* 1999;33:327-32.
- Schwartz PJ, Priori SG, Napolitano C. How really rare are rare diseases? The intriguing case of independent compound mutations in the long QT syndrome. *J Cardiovasc Electrophysiol* 2003;14:1120-1.
- Roden DM. Human genomics and its impact on arrhythmias. *Trends Cardiovasc Med* 2004;14:112-6.
- Taggart NW, Haglund CM, Tester DJ, Ackerman MJ. Diagnostic miscues in congenital long-QT syndrome. *Circulation* 2007;115:2613-20.
- Maron BJ. Sudden death in young athletes. *N Engl J Med* 2003;349:1064-75.
- Choi G, Kopplin LJ, Tester DJ, Will ML, Haglund CM, Ackerman MJ. Spectrum and frequency of cardiac channel defects in swimming-triggered arrhythmia syndromes. *Circulation* 2004;110:2119-24.
- Viskin S, Rosovski U, Sands AJ, et al. Inaccurate electrocardiographic interpretation of long QT: the majority of physicians cannot recognize a long QT when they see one. *Heart Rhythm* 2005;2:569-74.
- Schwartz PJ, Moss AJ, Vincent GM, Crampton RS. Diagnostic criteria for the long QT syndrome: an update. *Circulation* 1993;88:782-4.
- Hofman N, Wilde AA, Käb S, et al. Diagnostic criteria for congenital long QT syndrome in the era of molecular genetics: do we need a scoring system? *Eur Heart J* 2007;28:575-80.
- Yang P, Kanki H, Drolet B, et al. Allelic variants in long QT disease genes in patients with drug-associated torsades de pointes. *Circulation* 2002;105:1943-8.

33. Paulussen AD, Gilissen RA, Armstrong M, et al. Genetic variations of KCNQ1, KCNH2, SCN5A, KCNE1, and KCNE2 in drug-induced long QT syndrome patients. *J Mol Med* 2004;82:182-8.
34. Lehtonen A, Fodstad H, Laitinen-Forsblom P, Toivonen L, Kontula K, Swan H. Further evidence of inherited long QT syndrome gene mutations in antiarrhythmic drug-associated torsades de pointes. *Heart Rhythm* 2007;4:603-7.
35. Arnestad M, Crotti L, Rognum TO, et al. Prevalence of long-QT syndrome gene variants in sudden infant death syndrome. *Circulation* 2007;115:361-7.
36. Shimizu W, Noda T, Takaki H, et al. Epinephrine unmasks latent mutation carriers with LQT1 form of congenital long-QT syndrome. *J Am Coll Cardiol* 2003;41:633-42.
37. Zhang L, Timothy KW, Vincent GM, et al. Spectrum of ST-T-wave patterns and repolarization parameters in congenital long-QT syndrome: ECG findings identify genotypes. *Circulation* 2000;102:2849-55.
38. Tester DJ, Will ML, Haglund CM, Ackerman MJ. Effect of clinical phenotype on yield of long QT syndrome genetic testing. *J Am Coll Cardiol* 2006;47:764-8.
39. Van Langen I, Birnie E, Alders M, Jongbloed RJ, Le Marec H, Wilde AA. The use of genotype-phenotype correlations in mutation analysis for the long QT syndrome. *J Med Genet* 2003;40:141-5.
40. Tester DJ, Cronk LB, Carr JL, et al. Allelic dropout in long QT syndrome genetic testing: a possible mechanism underlying false-negative results. *Heart Rhythm* 2006;3:815-21.
41. Kimbrough J, Moss AJ, Zareba W, et al. Clinical implications for affected parents and siblings of probands with long-QT syndrome. *Circulation* 2001;104:557-62.
42. Hobbs JB, Peterson DR, Moss AJ, et al. Risk of aborted cardiac arrest or sudden cardiac death during adolescence in the long-QT syndrome. *JAMA* 2006;296:1249-54.
43. Sauer AJ, Moss AJ, McNitt S, et al. Long QT syndrome in adults. *J Am Coll Cardiol* 2007;49:329-37.
44. Brink PA, Crotti L, Corfield V, et al. Phenotypic variability and unusual clinical severity of congenital long-QT syndrome in a founder population. *Circulation* 2005;112:2602-10.
45. Schwartz PJ, Priori SG, Cerrone M, et al. Left cardiac sympathetic denervation in the management of high-risk patients affected by the long-QT syndrome. *Circulation* 2004;109:1826-33.
46. Benhorin J, Taub R, Goldmit M, et al. Effects of flecainide in patients with new SCN5A mutation: mutation-specific therapy for long-QT syndrome? *Circulation* 2000;101:1698-706.
47. Moss AJ, Windle JR, Hall WJ, et al. Safety and efficacy of flecainide in subjects with long QT-3 syndrome (DeltaKPQ mutation): a randomized, double-blind, placebo-controlled clinical trial. *Ann Noninvasive Electrocardiol* 2005;10:Suppl:59-66.
48. Priori SG, Napolitano C, Schwartz PJ, Bloise R, Crotti L, Ronchetti E. The elusive link between LQT3 and Brugada syndrome: the role of flecainide challenge. *Circulation* 2000;102:945-7.
49. Makita N, Behr E, Shimizu W, et al. Overlap between LQT3 and Brugada syndrome: clinical features in a common mutation and underlying biophysical mechanisms. *Circulation* 2007;116:Suppl II:II-490. abstract.
50. Moss AJ, Zareba W, Hall WJ, et al. Effectiveness and limitations of beta-blocker therapy in congenital long-QT syndrome. *Circulation* 2000;101:616-23.
51. Daubert JP, Zareba W, Rosero SZ, Budzikowski A, Robinson JL, Moss AJ. Role of implantable cardioverter defibrillator therapy in patients with long QT syndrome. *Am Heart J* 2007;153:Suppl:53-8.
52. Zipes DP, Camm AJ, Borggrefe M, et al. ACC/AHA/ESC 2006 guidelines for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force and the European Society of Cardiology Committee for Practice Guidelines (writing committee to develop Guidelines for Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 2006;114(10):e385-e484.

Copyright © 2008 Massachusetts Medical Society.

COLLECTIONS OF ARTICLES ON THE JOURNAL'S WEB SITE

The *Journal's* Web site (www.nejm.org) sorts published articles into more than 50 distinct clinical collections, which can be used as convenient entry points to clinical content. In each collection, articles are cited in reverse chronologic order, with the most recent first.